

Human TL1A Protein; His Tag


Product Information

| | |
|------------------------|---|
| Product Name | Human TL1A Protein; His Tag |
| Storage temp. | Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Catalog# / Size | GM-84079RP-100 / 100 μg GM-84079RP-1000 / 1 mg |

Protein Information

| | |
|---------------------------|---|
| Alternative Names | VEGI, TNFSF15 |
| Source | Human TL1A Protein; His Tag (GM-84079RP) is expressed from human 293 cells (HEK-293). It contains AA Leu 72 - Leu 251 (Accession #O95150-1). This protein carries a His tag at the N-terminus. |
| Purity | > 90% as determined by SDS-PAGE |
| Endotoxin | < 1 EU/ μg , determined by LAL gel clotting assay |
| Predicted Mol Mass | 22.4 KDa |
| Formulation | Supplied as a 0.2 μm filtered solution of PBS, pH7.4. |
| Description | Tumor necrosis factor (TNF)-like cytokine 1A (TL1A) is a member of the TNF superfamily of proteins (TNFSF15), which signals through association with death domain receptor 3 (DR3). Decoy receptor 3 (DcR3) competes with DR3 for TL1A binding and inhibits functional signaling. These proteins are significantly upregulated in inflamed intestinal tissues, and their pathogenetic importance for inflammatory bowel disease (IBD) is suggested by accumulating evidence. TL1A/DR3 induce costimulatory signals to activated lymphocytes, including the gut-specific populations of CD4+CD161+ and CD4+CCR9+ cells, affecting all major effector pathways and inducing the mucosal upregulation of Th1, Th2, and Th17 factors. |


SDS-PAGE



On SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90%.


Bioactivity-ELISA

Human TL1A Protein; His Tag, ELISA 0.5 µg of Human TL1A Protein; His Tag per well




Human TL1A Protein; His Tag (Catalog # GM-84079RP) was immobilized at 5 µg/ml (100 µL/well). Increasing concentrations of Anti-H_TNFSF15(TL1A) hIgG1 Antibody (Tulisokibart、PRA-023) (Catalog # GM-58915AB) were added.

Bioactivity CELL BASE



Human TL1A Protein; His Tag (Catalog # GM-84079RP) was added into H_TNFRSF25(DR3) Reporter Jurkat Cell Line (Catalog # GM-C30290), and stimulates TL1A/DR3 signals.



| | |
|------------------|---------------------------------|
| | H_TNFRSF25(DR3) Reporter Jurkat |
| IC ₅₀ | 0.05321 |

Neutralization assay shows that the stimulate effect of 10 ng/mL Human TL1A Protein; His Tag (Catalog # GM-84079RP) was inhibited by increasing concentration of Anti-H_TNFSF15(TL1A) hIgG1 Antibody (Tulisokibart、PRA-023)(Catalog # GM-58915AB).