

# Product Sheet

## H\_ALPPL2(ALPG) HEK-293 Cell Line

Catalog number: GM-C26399

Version 3.3.1.250806

<b>Description</b>	H_ALPPL2(ALPG) HEK-293 Cell Line is a clonal stable HEK-293 cell line that constitutively expresses the human ALPPL2(ALPG) gene, constructed using lentiviral technology.
<b>Quantity</b>	5E6 Cells per vial, 1 mL
<b>Product Format</b>	1 vial of frozen cells
<b>Shipping</b>	Shipped on dry ice
<b>Storage Conditions</b>	Liquid nitrogen immediately upon receipt
<b>Target</b>	Human_ALPPL2(ALPG)
<b>Gene ID/Uniprot ID</b>	P10696
<b>Host Cell</b>	HEK-293
<b>Recovery Medium</b>	DMEM+10% FBS+1% P.S
<b>Growth medium</b>	DMEM+10% FBS+1% P.S+0.75 µg/mL Puromycin
<b>Note</b>	None
<b>Freezing Medium</b>	90% FBS+10% DMSO
<b>Growth properties</b>	Adherent
<b>Growth Conditions</b>	37°C, 5% CO <sub>2</sub>
<b>Mycoplasma Testing</b>	The cell line has been screened to confirm the absence of Mycoplasma species.
<b>Safety considerations</b>	Biosafety Level 2
<b>Note</b>	It is recommended to expand the cell culture and store a minimum of 10 vials at an early passage for potential future use.

## Materials

Reagent	Manufacturer/Catalogue No.
DMEM	Gibco/C11995500BT
Fetal Bovine Serum	ExCell/FSP500
Pen/Strep	Thermo/15140-122
Puromycin	Genomeditech/ <a href="#">GM-040401</a>
Anti-ALPPL2 mIgG2a antibody(SGN-ALPV)	Genomeditech/ <a href="#">GM-58773AB</a>

## Figures




Figure 1 | H\_ALPPL2(ALPG) HEK-293 Cell Line (Cat. GM-C26399) Was determined by flow cytometry using Anti-ALPPL2 mIgG2a antibody(SGN-ALPV) (Cat. [GM-58773AB](#)).




Figure 2 | The passage stability of the H\_ALPPL2(ALPG) HEK-293 Cell Line (Cat. GM-C26399) was determined by flow cytometry using Anti-ALPPL2 mIgG2a antibody(SGN-ALPV) (Cat. [GM-58773AB](#)).

## Cell Recovery

Recovery Medium: DMEM+10% FBS+1% P.S

To insure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

- Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 - 3 minutes).
- Remove the vial from the water bath as soon as the contents are thawed, and decontaminate by dipping in or spraying with 70% ethanol. All of the operations from this point on should be carried out under strict aseptic conditions.
- Transfer the vial contents to a centrifuge tube containing 5.0 mL complete culture medium and spin at approximately 176 x g for 5 minutes. Discard supernatant.
- Resuspend cell pellet with the recommended recovery medium. And dispense into appropriate culture dishes.
- Incubate the culture at 37°C in a suitable incubator. A 5% CO<sub>2</sub> in air atmosphere is recommended if using the medium described on this product sheet.

## Cell Freezing

Freezing Medium: 90% FBS+10% DMSO

- Centrifuge at 176 x g for 3 minutes to collect cells.
- Resuspend the cells in pre-cooled freezing medium and adjust the cell density to 5E6 cells/mL.
- Aliquot 1 mL into each vial.
- Place the vial in a controlled-rate freezing container and store at -80°C for at least 1 day, then transfer to liquid nitrogen as soon as possible.

## Cell passage

Growth medium: DMEM+10% FBS+1% P.S+0.75 µg/mL Puromycin

For the first 1 to 2 passages post-resuscitation, use the recovery medium. Once the cells have stabilized, switch to a growth medium.

- Subculturing is necessary when the cell density reaches 80%. It is recommended to perform subculturing at a ratio of 1:3 to 1:4 every 2-3 days. Ensure that the density does not exceed 80%, as overcrowding can lead to reduced viability due to compression.
- Remove and discard culture medium.
- Briefly rinse the cell layer with PBS to remove all traces of serum that contains trypsin inhibitor.
- Add 1.0 mL of 0.25% (w/v) Trypsin-EDTA solution to dish and observe cells under an inverted microscope until cell layer is dispersed (usually within 30 to 60 seconds at 37°C).
- Note: To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
- Add 2.0 mL of growth medium to mix well and aspirate cells by gently pipetting.

- g) After centrifugation, resuspend the pellet and add appropriate aliquots of the cell suspension to new culture vessels.
- h) Incubate cultures at 37°C.

**Subcultivation Ratio: A subcultivation ratio of 1:3 - 1:4 is recommended**

**Medium Renewal: Every 2 to 3 days**

## Notes

- a) Upon initial thawing, a higher number of dead cells is observed, which is a normal phenomenon. Significant improvement is seen after adaptation. Once the cells reach a stable state, the number of dead cells decreases after subculturing and the cell growth rate becomes stable.
- b) Ensure that the cell density does not exceed 80%, as overcrowding may lead to reduced viability due to compression.

## Sequence

ALPPL2(ALPG) [P10696](#)

MQGPWVLLLLGLRLQLSLGIIPVEEENPDFWNRQAAEALGAAKKLQPAQTAACKNLIIFLGDGMGVSTVTAAR  
ILKGQKKDKLGPETFLAMDRFPYVALSKTYSVDKHVPDSGATATAYLCGVKGNFQTIGLSAAARFNQCNTT  
RGNEVISVMNRAKKAGKSVGVTTRVQHASPAGAYAHTVNRNWYSDADVPASARQEGCQDIATQLISNM  
DIDVILGGGRKYMFPMTGPDPEYPDDYSQGGTRLDGKNLVQEWLAKHQGARYVWNRTELLQASLDPSVTH  
LMGLFEPGDMKYEIHRDSTLDPSEMEMTEAALLLSRNPRGFFLFVEGGRIDHGHESRAYRALTIMFDD  
AIERAGQLTSEEDTSLVTADHSHVFSFGGYPLRGSSIFGLAPGKARDRKAYTVLLYGNGPGYVLKDGARPD  
VTESESGSPEYRQQSAVPLDGETHAGEDVAVFARGPQAHLVHGVQEQTFAHVMAFAACLEPYTACDLAPR  
AGTTDAAHPGPSVVPALLPLLAGTLLLLGTATAP

## Related Products

ALPPL2 ALPP	
<a href="#">Cynomolgus_ALPP HEK-293 Cell Line</a>	<a href="#">H_ALPI HEK-293 Cell Line</a>
<a href="#">H_ALPL HEK-293 Cell Line</a>	<a href="#">H_ALPP CHO-K1 Cell Line</a>
<a href="#">H_ALPP CT26 Cell Line</a>	<a href="#">H_ALPP HEK-293 Cell Line</a>
<a href="#">H_ALPP LLC1 Cell Line</a>	<a href="#">H_ALPP MC38 Cell Line</a>
<a href="#">H_ALPPL2(ALPG) CHO-K1 Cell Line</a>	<a href="#">H_ALPPL2(ALPG) CT26 Cell Line</a>
<a href="#">H_ALPPL2(ALPG) LLC1 Cell Line</a>	<a href="#">H_ALPPL2(ALPG) MC38 Cell Line</a>
<a href="#">Anti-ALPPL2 hIgG1 antibody(SGN-ALPV)</a>	<a href="#">Anti-ALPPL2 mIgG2a antibody(SGN-ALPV)</a>
<a href="#">Anti-H_ALPPL2 hIgG1 Reference Antibody(h12F3)</a>	<a href="#">Anti-ALPPL2-MMAE ADC(Dar4)[SGN-ALPV]</a>
<a href="#">Biotinylated Cynomolgus ALPI Protein; His-Avi Tag</a>	<a href="#">Biotinylated Cynomolgus ALPL Protein; His-Avi Tag</a>
<a href="#">Biotinylated Cynomolgus ALPP Protein; His-Avi Tag</a>	<a href="#">Biotinylated Human ALPI Protein; His-Avi Tag</a>
<a href="#">Biotinylated Human ALPL Protein; His-Avi Tag</a>	<a href="#">Biotinylated Human ALPP Protein; His-Avi Tag</a>
<a href="#">Cynomolgus ALPI Protein; His Tag</a>	<a href="#">Cynomolgus ALPL Protein; His Tag</a>
<a href="#">Human ALPI Protein; His Tag</a>	<a href="#">Human ALPL Protein; His Tag</a>
<a href="#">Human ALPP Protein; His Tag</a>	<a href="#">Human ALPPL2 Protein; His Tag</a>

ADC Related Product	
Anti-DXD Mouse IgG1 Antibody (23E21C5)	Anti-DXD Mouse IgG1 Antibody (4A5A12)
Anti-Dxd Mouse IgG2a Antibody (17D6A4)	Anti-Eribulin Mouse IgG2a Antibody (10F8G4)
Anti-MMAE Mouse IgG1 Antibody (11C10E3)	Anti-MMAE Mouse IgG2a Antibody (17A1K11)
Anti-MMAE Mouse IgG2a Antibody (8F6A3)	Mouse anti Human IgG1-MMAE(Dar4)
Human IgG1 Isotype-DXD (Dar8)	Human IgG1 Isotype-Eribulin (Dar4)
Human IgG1 Isotype-MMAE (Dar4)	
Recombinant DT3C Protein	

## License Agreement:

**By purchasing and using this cell line product, the user voluntarily agrees to accept and abide by the following policies:**

- This cell line product is restricted to research use only and shall not be used for any commercial purposes.
- This product is strictly prohibited from being used in the diagnosis or treatment of human or animal diseases, and shall not be directly used in experiments involving humans.
- Users and their contractors engaged for their benefit may use this material and its derivatives only within the agreed research scope; modification of the material is not permitted, nor may it be distributed, sold, transferred, or otherwise provided to any other entity (including affiliates).
- If use beyond the above scope is required, prior written permission from Genomeditech (Shanghai) Co.,Ltd. must be obtained. For details, please contact Genomeditech (Shanghai) Co.,Ltd.